


# Make sure your child sees, hears and listens

## Developmental Chart




NOTE: To use this chart, keep a pencil vertically or the age of the child. All milestones falling to the left of the pencil should have been achieved by the child.

Based on BSID Baroda norms & Trivandrum Developmental screening chart (TDSC)

— This represents normal range

Doctor's Name :

Dr. Prasanna



*Good Health and Good Sense are  
two life's greatest blessings*

Immunization and Health Card

*Shankar's Hospital*





# My Health Record

Dear Dad and Mom, This is my immunization and health record diary. This will help you to understand my health needs and guidance on my care. Please read the information carefully and always carry it every time you take me to our doctor.

Your cutie.

**ZAHRA**

## My details:

My name: ZB Aminah

I am a boy  girl

Mom's name: Aminah

Dad's name: \_\_\_\_\_

Home address: \_\_\_\_\_

Contact number: \_\_\_\_\_

## Birth details:

Date and time of birth: 6/8/16 at 8:54am

Place of birth: Dharmalashmi Hospital

Delivery: Normal / vacuum/ forceps/ caesarean

Delivered by (Obstetrician): \_\_\_\_\_

Birth weight: 2.8 kg Length at birth: \_\_\_\_\_

H.C: \_\_\_\_\_

Blood Group: B

Remarks: \_\_\_\_\_

## General instructions:

1. BCG vaccination causes a small swelling which will heal slowly by itself forming a scar and no medicines should be applied on it.
2. Children can be breast fed soon after giving polio drops.
3. If vaccination was discontinued or delayed, it could be continued from where it was stopped.
4. Even weak children should take vaccination for protection.
5. Children with mild fever, cold, cough or diarrhea can take all the vaccines.

# Vaccination Record

Age	Vaccine	Due date	Given date	Sign of Dr.
Birth	BCG Oral Polio Vaccine zero dose Hepatitis B-I dose		6/8/16	
6 weeks	DPT - I dose Hepatitis B - II dose Hib vaccine I dose OPV - I dose BCG (If not given at Birth)		23/9/16 5-10-16	
10 weeks	DPT - II dose Hib vaccine II dose OPV - II dose		22/11/16 22/2/2017	
14 weeks	DPT - III dose Hib vaccine III dose OPV - III dose		22/3/17 19-4-17	
18 months	Measles Hepatitis B III dose MMR		20/12/2017 16/10/19	
18 months	DPT - I Booster OPV - I Booster Hib Vaccine - Booster dose		16/10/19	
5 - 6 years	DPT - II Booster			
10 years	TT Booster dose			
16 years	TT Booster dose			
Not under national immunization schedule.				
6th Weeks to 6 Month	Vaccine against 1. Rota Virus - 2 doses 2. Pneumococci - 3 doses			
1 Year	Chicken Pox Vaccine Hepatitis A Vaccine			
18 Months	Hepatitis A 2nd dose (after 6 months of 1 dose)			

## Note:

1. A dose of Hepatitis B Vaccine within 12 hours of birth is essential to prevent mother to child transmission.
2. If Hepatitis B Vaccine is given at birth, remaining doses can be given at 10, 14 weeks or alternatively 6 & 14 weeks.
3. BCG, OPV and Hepatitis B vaccines, when missed at birth can be started at the completion of 6 weeks.
4. For Typhoid immunization, earliest age recommended: Whole cell Vaccines at 6 months, Vi antigen at 2 years and Ty21a vaccine at 6 years. Booster dose every 3 years.
5. In addition to routine OPV doses the recommended 'Pulse OPV dose' are also mandatory during PPI campaigns.
6. 2 doses of Influenza Vaccine are given when immunizing 1st time.
8. DTP vaccine includes DTPw or DTPa.