Vaccine \ Age	Birth	2 Months	4 Months	6 Months	12 Months	15 Months	18 Months	4 - 6 Years	13 - 14 Years
BCG	BCG								
Hepatits B	HBV 0								
A Penta (DTP+Hib+HBV)		Penta 1		Penta 2		C Penta 3		D (Booster)	Tdap
B Hexa (DTaP+ Hib+HBV+IPV)			Hexa						
Rotavirus		Rota 1	Rota 2						
Oral Polio Vaccine		OPV 1	OPV 2	OPV 3			OPV (Booster)	OPV (Booster)	
Pneumococcal Conjugate		PCV 1	PCV 2	PCV 3		PCV 4			
Measles, Mumps, Rubella					MMR 1		MMR 2		
Varicella					Varicella 1			Varicella 2	
Hepatitis A					Hep A 1		Hep A 2		
Influenza				Influenza (Yearly)					
Meningococcal ACYW135									For adults & children > 2 years

Your Child's Immunization Schedule

A. Penta: DTP (Diphtheria, Tetanus, Pertusis) + Hib (Haemophilus influenza type B) + HBV (Hepatitis B)

B. Hexa: DTaP (Diphtheria, Tetanus, acellular Pertusis) +Hib (Haemophilus influenza type B) + HBV (Hepatitis B) + IPV (Inactivated Polio)

C. Tetra: (DTaP+Hib) (Diphtheria, Tetanus, acellular Pertusis) +Hib (Haemophilus influenza type B)

D. DTaP: (Booster) (Diphtheria, Tetanus, acellular Pertusis)

E. Tdap: (Tetanus, diphtheria, acellular pertusis)

Vaccines recommended for routine administration to children

Highly recommended for all from 6 months yearly

Meningococcal meningitis for Hajj, Umrah & Endemic countries and for children after 2 years.

Following immunization inform the doctor/nurse if your Child is suffering from:

- Epileptic fits, seizures or convulsions
- Any untoward reaction from the previous vaccinations
- Severe cough/colds with fever and not feeling well in anyway

Common Immunization Reactions:
Fever, local swelling, redness and pain at the injection site

Home Care Advice for Immunization Reactions
Local Reaction at injection Site:
Cold Pack : 20 minutes each hour as needed
Fever: Give Acetaminophen or Ibuprofen as needed.
Localized Hives: Apply 1 % hydocortisone cream OTC once or twice.

Next Vaccination	
Date	Vaccine

For more information Please call us at:
4408-2444
Email : alahlireception@qatargas.com.qa