

Name BABY TUBA. Sex \_\_\_\_\_

Date of Birth \_\_\_\_\_ Time \_\_\_\_\_

Brith History \_\_\_\_\_

Significant Information \_\_\_\_\_

### IMMUNISATION RECORD

AGE	VACCINE	DUE ON	GIVEN ON
Birth	B.C.G.		
	Oral Polio - 1st Dose		
6 Weeks	DPT - 2nd Dose		
	Oral Polio - 2nd Dose		
	Hepatitis B 1st Dose		
10 Weeks	DPT -2nd Dose		
	Oral Polio - 3rd Dose		
	Hepatitis B 2nd Dose		
14 Weeks	DPT - 3rd Dose		
	Oral Polio 4th Dose		
18 Weeks	Oral Polio- 5th Dose		
6 Months	Hepatitis B 3rd Dose		
9 Months	Measles		
15 Months	MMR		
18-24 Months	DPT- Polio 1st Booster Dose		
2- 3 Years	Typhoid Vaccine*		
5 Years	DPT - Polio 2nd Booster Dose		
	Hepatitis B - Booster Dose		
6-8 Years	Typhoid Vaccine		
10 Years	T.T. (3rd Booster Dose)		
15-16 Years	T.T. (Tetanus) - 4th Booster Dose		

#### NEW VACCIENS H. INFUENZA B. VACCINE

	Due on	Given on
1st Dose		
IIInd Dose		
IIIrd Dose		
Booster Dose		

#### HEPATITIS - A - VACCINE

	Due on	Given on
1st Dose		
IIInd Dose		

#### IPV

	Due on	Given on
1st Dose		
IIInd Dose		
IIIrd Dose		
Booster Dose		

#### ROTAVIRUS VACCINE

	Due on	Given on
1st Dose		
IIInd Dose		

#### PLEASE NOTE

- Inform your doctor if child has significant reaction to the last dose.
- D.P.T. : There may be fever, redness, swelling at site of injection for two days, child may develop nodule which will disappear after few weeks.
- H. Influenza B Vaccine is for prevention of Pneumania and Meningitis.
- Do not ask for medicines on Phone.
- For Appointment on Sunday  
(Call at 9.00 a.m. on 26658514)

#### ADDITIONAL VACCINES

VAXIGRIP			
PREVENAR VACCINE			
→ CHICKEN POX → 2 / 7/2017 / 7/2017			