

AGE	VACCINE (dose)	GIVEN ON	SIGNATURE
Birth	BCG Hep B Vac		
2 Months (1st Dose)	DPT + Hib AUSTRA 322A OPV-eSas Hep B Vac A15CA391A Pneumococcal - 9V4932 (Merk) DPT + Hib A15CA391A OPV - t5510 - 91-11 Hep B Vac A15CA391A Pneumococcal - ES4095	22/11/10 22/11	Susan
4 Months (2nd Dose)	DPT + Hib A15CA391A OPV - t5510 - 91-11 Hep B Vac A15CA391A Pneumococcal - ES4095	22/11	Susan
6 Months (3rd Dose)	DPT + Hib A15CA391A OPV - t5510 - 91-11 Hep B Vac A15CA391A Pneumococcal - ES4095	13/11/11	Susan
12 Months	MMR 1 - A WJR B990AA Chickenpox - A T003 G59A Narcellig) Hep A - A HNR B3484BF	17/10/2011	Nedie
18 Months (1st Dose)	MMR - A WJR B990AA OPV - A OPV B871AD HEPA - A HAV B3484BF	22/4/11	CP
4-6 Years (1st Booster)	DT OPV MMR 2		
11-12 Years	Tetanus OPV		

TETRA - G 0169 } 11/3/12 CP
PCV - F 22728 }

Before the immunization tell your doctor if your child:

- Has ever had fits or convulsions
- Had a bad reaction to the last dose
- Is unwell in anyway

Current Immunization Schedule in State of Qatar 2005




"Common adverse effect!"

- Pain
- Redness and Swelling at Injection site
- Fever

"Treatment"

- Give paracetamol every 6 to 8 hours as needed
- Put warm compress on injection site
- If worried about child call doctor



Breast Feeding:

- Breast milk contains a balance of nutrients
- Breast milk is easy for baby to digest
- Breast milk contains antibodies that protect your baby
- Breast milk actually changes to keep up with baby.
- Breast - feeding creates mother child bonding.

Weaning is the introduction of solid food... How to Start ?

The best way to decide how and what food to give is to follow baby's development.

- Stage 1 - First taste - smooth foods around 6 - 7 months
 - Stage 2 - Learn to chew - soft lumps 8 - 9 months
 - Stage 3 - Self feeding - finger food, lumps 9 - 12 months
 - Stage 4 - Family diet with modifications after 12 months.
- ✗ Avoid giving raw vegetables, raisins, hotdogs or meat chunks, nuts or hard candies before 2 years.
- ✗ No cow's milk for the 1st year.