

B

**HEALTH RECORD**

Child's Name: Abdul Ahad Qamar S/O Qamar Zaman

Medical Record No: 577-24-09

Age/Date of Birth: 13/2/17

Place of Birth: Atanku Born Hyderabad

Family Name: Qamar

Address: Kalhora Colony, Hyderabad

Telephone: 4923322058633

**Important**  
Please keep this vaccination booklet in a safe place and bring it along with you on every visit.

**THE AGA KHAN UNIVERSITY HOSPITAL AND AFFILIATED CLINICAL SITES  
IMMUNIZATION SCHEDULE FOR CHILDREN**

Age	Name of Vaccines	Dose	Route
At Birth	1. BCG	0.05 ml (0-3 mo) 0.1 ml (> 3 mo)	Intradermal (preferred deltoid)
	2. Polio Virus (OPV)	2 drops	PO
6 Weeks	1. Diphtheria-Pertussis-Tetanus-Hepatitis B-Hemophilus Influenza type b	0.5 ml	IM
	2. Polio Virus (OPV)	2 drops	PO
	3. Pneumococcal-Conjugate Vaccine	0.5 ml	IM
10 Weeks	1. Diphtheria-Pertussis-Tetanus-Hepatitis B-Hemophilus Influenza type b	0.5 ml	IM
	2. Polio Virus (OPV)	2 drops	PO
	3. Pneumococcal Conjugate Vaccine	0.5 ml	IM
	4. Rotavirus*	1 ml	PO
14 Weeks	1. Diphtheria-Pertussis-Tetanus-Hepatitis B-Hemophilus Influenza type b	0.5 ml	IM
	2. Polio Virus (OPV)	2 drops	PO
	3. Pneumococcal Conjugate Vaccine	0.5 ml	IM
	4. Rotavirus*	1 ml	PO
9 Months	1. Measles / MMR	0.5 ml	SC <i>Eniscent</i>
12 Months	1. Hepatitis A*	0.5 ml	IM
	2. Measles/Measles-Mumps-Rubella (MMR) or MMR-Varicella*	0.5 ml	SC
15 Months	1. Measles-Mumps-Rubella or MMR-Varicella**	0.5 ml	SC
	2. Pneumococcal Conjugate Vaccine*	0.5 ml	IM
18 Months	1. Diphtheria-Pertussis-Tetanus-Hepatitis B-Hemophilus Influenza type b	0.5 ml	IM
	2. Polio Virus (OPV)	2 drops	PO
	3. Hepatitis A*	0.5 ml	IM
> 2 Years	1. Typhoid*	0.5 ml	IM
	2. Varicella (if not given earlier)*	0.5 ml	SC
	3. Meningococcal Conjugate Vaccine*		IM
4-5 Years	1. Diphtheria-Tetanus-Pertussis-Hepatitis B Hib**	0.5 ml	IM
	2. Measles-Mumps-Rubella or MMR-Varicella**	0.5 ml	SC
	3. Polio Virus (OPV)	2 drops	PO

a. MMR-Varicella should be given if available. If MMRV not available given MMR at 15 months and Varicella can be given at 2 years and 4-5 years as separate injections.  
b. Tetanus toxoid 0.5 ml IM is indicated for children between 10-15 years of age.  
\* These vaccines are not currently supported by the National Expanded Programme for Immunisations (EPI), Pakistan, because of expense issues, but are strongly recommended by the Department of Paediatrics and Child Health, AKU.  
PO = Per Oral - IM = Intramuscular - SC = Subcutaneous

Age	Name of Vaccines	Given on Date / Time	Next Dose Date	Nurse's Name with Designation
	Bacuta	15/1/17		Jhanna
	Penta OPV Polio	2/1/17	1-3-17	Amir
	Penta OPV Polio	4/1/17	3/4/17	dur green
	Penta OPV Polio	25/1/17	5/1/17	dur. wmm
	Measles	24/1/17	13/10/17	KA
	Polio Hib Measles	4/3/17	11/1/18	KA
	PCV13	20.4.18	13/4/18	A2
	Hib	13/1/18	7/1/18	A2
	Measles	30/1/18	15/1/19	KA
	DPH Polio	3/1/21	13/1/21	Boz

Mon to sat  
9am to 5pm