

M. S. SANTO CHRIS

GOKULAM HOSPITAL

CHITHAMBARA VEETHI,
Kottar, Nagercoil

S. SANTO CHRIS



S SANKI TO CRIS.

Date of Birth _____ Birth Weight _____ Length _____ Head Circumference _____

Delivery: Normal / Vacuum / Forceps / Caesarian _____ Age: _____ Sex: _____

Remarks _____

Vaccination Record

Age	Vaccine	Due date	Given date	Comments/ Peel-off sticker	
Birth	BCG Vaccine	}	30/12/11	114	
	OPV 0 dose + Hepatitis B1				
6 weeks	DTwP 1 /DTap 1				
	OPV 1* / OPV 1 + IPV 1				
	Hib 1				
	Hepatitis B2				
	Rotavirus 1 **	16	2	15	
	PCV 1				
10 weeks	DTwP 2 /DTap 2				
	OPV 2* / OPV 2 + IPV 2	16	2	15	
	Hib 2				
	Rotavirus 2 **				
	PCV 2				
14 weeks	DTwP 3 /DTap 3				
	OPV 3* / OPV 3 + IPV 3				
	Hib 3	18	4	15	
	Hepatitis B3				
	Rotavirus 3 **				
PCV 3					
10 Months	Measles		21	10	15
12 Months	Hepatitis A 1				
15 Months	MMR 1		17	3	16
16 Months	Varicella				
	PCV Booster				
18 Months	DTwP B1 / DTap B1		23	6	16
	OPV 4* / OPV 4 + IPV B1				
	Hib B1				
< 2 Years	Hepatitis A2				
2 Years	Typhoid 1*		18	1	17
5 Years	DTwP B ₂ / DTap B ₂				
	OPV 5				
	MMR 2*				
	Typhoid 2	24	1	19	
	Varicella 2 **				
10 to 12 Years	Tdap / Td*				
	HPV ^				

Don't have
 + 15w

 Don't have
 15w

 Don't have
 15w

 Don't have
 15w

Parents to Note

Prior to immunization, inform your doctor if the child has had any significant reaction to the last dose. Immunization can be given in the presence of a minor illness.

After the immunization, reactions are usually mild

BCG

A nodule appears 3-4 weeks after BCG Vaccination. It may soften or ulcerate in 2-4 weeks. No application or formation is necessary. It heals, leaving a scar, indicating effective vaccination.

DTwP

There may be mild fever and pain, redness and swelling at the site of the injection. A small painless lump may remain for a few weeks. For fever and pain, paracetamol syrup/tablet may be given, the dose can be repeated every 4-6 hours if required. Please consult your Doctor for any other reaction.

Measles/MMR

A few children get fever 4 to 10 days after the vaccination. Paracetamol syrup/tablets may be given if required.

- Additional vaccines may be given under special circumstances
- Vaccines are usually safe. The benefits of vaccination outweigh any risk associated with the immunization
- All vaccines, through properly given may not give full protection from the diseases

The schedule may be modified by your doctor.

Dr M BAGAVATHY PERUMAL MD (HEAD)
 Reg. No : 58177
 Gokulam Hospital, Kottar

Note - 1, One dose of Hep B vaccine should be given within 12 hrs of birth to prevent mother to child transmission. 2. In addition to OPV doses in the schedule, recommended Pulse Polio doses are compulsory during Polio Ravivar. 3. If missed the schedule, ask your doctor about catch-up vaccination.